

## Documentation of Weed species from Sakti District, Chhattisgarh, India

Pooja Chaudhary<sup>1</sup>, Anshu Deep Khalkho<sup>2\*</sup>, Roopshikha Agarwal<sup>3</sup>, Madhulika Tigga<sup>4</sup>,  
Alok Toppo<sup>5</sup>, Deepak Kumar Soni<sup>6</sup>

<sup>1,2</sup>Department of Botany, Bharti Vishwavidyalaya, Durg, Chhattisgarh, India

<sup>3</sup>Department of Botany, Government J. Y. Chhattisgarh College, Raipur, Chhattisgarh, India

<sup>4</sup>Department of Botany, Government Bala Saheb Deshpande college Kunkuri, Jashpur, Chhattisgarh, India

<sup>5</sup>Department of Botany, Government College Batauli, Sarguja, Chhattisgarh, India

<sup>6</sup>Department of Life Science (Botany), Dr. C.V. Raman, Kota, Bilaspur, Chhattisgarh, India

\*Corresponding author email- [anshudeepxalxo18@gmail.com](mailto:anshudeepxalxo18@gmail.com)

DOI: [https://doi.org/10.63001/tbs.2026.v21.i03.S.I\(3\).pp259-265](https://doi.org/10.63001/tbs.2026.v21.i03.S.I(3).pp259-265)

### KEYWORDS

Weed diversity,  
agroecosystem, Sakti  
district, Chhattisgarh,  
floristic composition,  
weed management,  
biodiversity

**Received on:**

03-05-2026

**Accepted on:**

03-06-2026

**Published on:**

23-07-2026

### Abstract

The present study investigates the diversity and distribution of weed flora in the agricultural and nonagricultural landscapes of Sakti district, Chhattisgarh, India. Systematic field surveys were conducted across various locations representing both irrigated lowland and rainfed upland farming systems during 2023–2024. Quadrat sampling and monthly observations were employed to document seasonal variations and species composition. A total of 72 weed species belonging to multiple families were recorded, indicating rich floristic diversity in the study area. Dominant families included Asteraceae, Poaceae, Fabaceae, Amaranthaceae, and Cyperaceae, reflecting their high adaptability, reproductive efficiency, and ecological amplitude. The study highlights that weeds exhibit significant variation in distribution due to differences in soil moisture, cropping patterns, and anthropogenic disturbances. Several invasive and noxious species such as *Parthenium hysterophorus*, *Chromolaena odorata*, *Lantana camara*, *Cyperus rotundus*, and *Cynodon dactylon* were found to be widely distributed, posing serious threats to crop productivity and native biodiversity. In addition to their negative impacts, many weed species were found to possess medicinal properties, indicating their potential ethnobotanical importance. The findings emphasize that understanding weed diversity is crucial for developing effective and sustainable weed management strategies. Proper identification, monitoring, and ecological assessment of weeds can help minimize crop losses and enhance agricultural productivity while conserving biodiversity. This study provides baseline data for future ecological and agronomic research in the region.

### Introduction

The study focuses on the assessment of weed diversity in the agricultural and nonagricultural landscapes of the Sakti region. Weeds play a crucial role in agro ecosystems, influencing crop yield, soil health, and overall farm productivity. Understanding the diversity and

distribution of weeds in a specific region is essential for developing effective weed management strategies and promoting sustainable farming practices. Weeds are the unwanted plants that are the most underestimated crop pests. Weed is a plant growing where it is not desired (Chaudhary et al., 2025). Weeds are unsightly, useless, injurious

plants and are unwanted plants but all unwanted plants may not be true weeds. The beginning of weed production represents a serious constraint to worldwide agriculture but when left uncontrolled they can cause over 80% yield loss (Cousens and Mortimer, 1995). The weed flora of the agriculture field has a large ecological amplitude, so they multiply and flourish well even in different environmental conditions. Since weeds have unique potentialities for adaptation they survive almost in any environment. Weed plants are pioneer of land that increases the diversity of plants and agricultural land by utilizing the environmental potential. The diversity of plants increases by different cropping system and maximizes crop production. Study of weed biology was started at early 1930s. A study was performed by Pavlychenko (1937), and described the significance of weed and crop interaction. In northeast Bell et al. in 1962 published a series of "The Biology of Canadian Weeds", including more than 100 weed species to support the significance of weed in early 1970s (Bhowmik, 1997).

Many weeds has medicinal properties like like diuretic, dysentery, hepatoprotective enhances hair growth, sedative, antimelanogenic, hepatitis, respiratory analeptic, antihelmintic, asthma, hair loss, joint swelling, paralysis, baldness, itches, ulcer, bilious, cataract, rheumatism, cold, blood disorder, skin diseases, elephantiasis, fever, asthma, snake bite, influenza, kidney disease, night blindness, menstrual pain, larvicidal activity, hemorrhoids, ulcer, fever, nasal congestion, abdominal pain, nervous and heart disease, cough, malaria, worm, respiratory disease, tuberculosis, gastric disorder, headache, diarrhea, toothache, anxiolytic, hepatoprotective, antiulcer, anticancer, antimicrobial, antinociceptive, hypolipemic, abortifacient, chicken pox, bronchitis, intestinal helminthiasis, antiperiodic

(Chouhan et al., 2025; Chaudhary and Khalkho 2025; Prakash and Khalkho 2025)

## Materials and Methods

The present study was conducted in selected agricultural and nonagricultural fields at various locations of Sakti district, Chhattisgarh, India. The district is characterized by diverse agricultural landscapes comprising both wet (irrigated lowland) and dry (rainfed upland) farming systems. Prior to field surveys, detailed information regarding land use patterns, cropping practices, and irrigation status was collected to avoid difficulties during sampling and data collection. Special attention was given to the classification of land types, soil moisture conditions, and accessibility of sites. This approach ensured systematic coverage of representative fields and improved the accuracy and reliability of observations recorded during the study. A systematic field survey was conducted through frequent visits and repeated quadrat sampling over a period of one and a half years (2023–2024) to record weed diversity and distribution. Surveys were carried out monthly to capture seasonal variations in weed occurrence (Chaudhary et al., 2025).

Clear and detailed photographs of weed plants were taken during field surveys using a mobile phone camera. Images were captured from both near and distant views to document the overall habit and growth form of each weed species. Special emphasis was given to photographing diagnostic characters such as leaves, stems, flowers, fruits, and inflorescences, which are essential for accurate identification. Photographs were taken from multiple angles and at different growth stages to avoid misidentification. These

visual records served as important supporting material for taxonomic confirmation and future reference during laboratory identification and documentation (Khalkho and Rai 2020).

Weeds were identified and confirmed from (8) and (10).

## Result and Discussion

| Scientific name                                   | Family        | Local name          | Place of collection  |
|---------------------------------------------------|---------------|---------------------|----------------------|
| <i>Acmella radicans</i> (Jacq.) R.K.Jansen        | Asteraceae    | White spot flower   | Kirari               |
| <i>Achyranthes aspera</i> L.                      | Amaranthaceae | Devil's horsewhip   | Dabhara nawapara     |
| <i>Acmella oleracea</i> (L.) R.K. Jansen          | Asteraceae    | Buzz buttons        | Gopalpur, chandrapur |
| <i>Ageratum conyzoides</i> L.                     | Asteraceae    | Chick weed          | Sukhada              |
| <i>Alternanthera sessilis</i> (L.) R.Br.          | Amaranthaceae | Sessile joyweed     | Chandrapur           |
| <i>Alternanthera philoxeroides</i> (mart.) griseb | Amaranthaceae | Alligator weed      | Gopalpur, chandrapur |
| <i>Alysicarpus monilifer</i> (L.) DC              | Fabaceae      | Moneyworts          | Kharsia              |
| <i>Argemone mexicana</i> L.                       | Papaveraceae  | Prickly poppy       | Saposh               |
| <i>Amaranthus spinosus</i> L.                     | Amaranthaceae | Spiny pigweed       | Kirari               |
| <i>Andrographis paniculata</i> (Burm.f.) Nees     | Acanthaceae   | Green chiretta      | Gobra                |
| <i>Antigonon leptopus</i> Hook. & Arn.            | Polygonaceae  | Queen's wreath      | Malkharauda          |
| <i>Barleria buxifolia</i> L.                      | Acanthaceae   | Box leaved Barleria | Sukhada              |
| <i>Bidens pilosa</i> L.                           | Amaranthaceae | Cobblers pegs       | Hasaud               |
| <i>Blumea lacera</i> (Burm.f.) DC.                | Asteraceae    | Kukraundha          | Ghughri chhapora     |
| <i>Calotropis procera</i> (Aiton) W.T. Aiton      | Apocynaceae   | Giant milkweed      | Jetha                |
| <i>Cardamine flexuosa</i> With.                   | Brassicaceae  | Wood bitter-cress   | Gopalpur, chandrapur |
| <i>Celosia argentea</i> L.                        | Amaranthaceae | Plumed cockscomb    | Saposh               |
| <i>Centella asiatica</i> (L.) Urban               | Apiaceae      | Indian pennywort    | Kirari               |
| <i>Chromolaena odorata</i> (L.) R.M.              | Asteraceae    | Jack in the Bush    | Gobra                |

| King & H. Rob.                                                       |                |                       |                         |
|----------------------------------------------------------------------|----------------|-----------------------|-------------------------|
| <i>Cleome viscosa</i> L.                                             | Cleomaceae     | Tick weed             | Hasaud                  |
| <i>Coldenia procumbens</i> L.                                        | Boraginaceae   | Creeoing coldenia     | Chandrapur              |
| <i>Coelachne simpliciuscula</i><br>(Wight & Arn. ex Steud.)<br>Munro | Poaceae        | Small Marsh Millet    | Sukhada                 |
| <i>Commelina benghalensis</i> L.                                     | Commelinaceae  | Tropical spiderwort   | Gopalpur,<br>chandrapur |
| <i>Croton bonplandianus</i> Baill.                                   | Euphorbiaceae  | Railway weed          | Ghughri chhapora        |
| <i>Cullen corylifolium</i> (L.) Medik                                | Fabaceae       | Bakuchi               | Bhudanpur               |
| <i>Cynodon dactylon</i> (L.) Pers.                                   | Poaceae        | Bermuda grass         | Gopalpur,<br>chandrapur |
| <i>Cyperus compressus</i> L.                                         | Cyperaceae     | Annual sedge          | Kharsia                 |
| <i>Cyperus difformis</i> L.                                          | Cyperaceae     | Rice sedge            | Kirari                  |
| <i>Cyperus eragrostis</i> Lam.                                       | Cyperaceae     | Umbrella sedge        | Malkharauda             |
| <i>Cyperus rotundus</i> L.                                           | Cyperaceae     | Nutgrass              | Hasaud                  |
| <i>Dactyloctenium aegyptium</i>                                      | Poaceae        | Crowfoot grass        | Sukhada                 |
| <i>Dentella repens</i> (L.) J.R.Forst                                | Rubiaceae      | Creeping lickstooop   | Ghughri chhapora        |
| <i>Digitaria sanguinalis</i> (L.)                                    | Poaceae        | Hairy crabgrass       | Jetha                   |
| <i>Echinochloa colona</i> (L.) Link                                  | Paniceae       | Jungle rice           | Saposh                  |
| <i>Eclipta prostrata</i> (L.) L.                                     | Asteraceae     | False Daisy           | Bhudanpur               |
| <i>Eleusine indica</i> (L.) Gaetrn.                                  | Poaceae        | Wiregrass             | Kharsia                 |
| <i>Euphorbia hirta</i> L.                                            | Euphorbiaceae  | Snakeweed             | Chandrapur              |
| <i>Euphorbia nutans</i> Lag.                                         | Euphorbiaceae  | Eyebane               | Gopalpur,<br>chandrapur |
| <i>Euphorbia prostrata</i> Aiton.                                    | Euphorbiaceae  | Dodhak                | Gobra                   |
| <i>Evolvulus alsinoides</i> (L.) L.                                  | Convolvulaceae | Dwarf morning glory   | Kirari                  |
| <i>Grangea maderaspatana</i> (L.) L.<br>Poir                         | Asteraceae     | Madras carpet         | Malkharauda             |
| <i>Grona triflora</i> (L.) H. Ohashi &<br>K. Ohashi                  | Fabaceae       | Creeping tick trefoil | Ghughri chhapora        |
| <i>Hygrophila auriculata</i>                                         | Acanthaceae    | Gokulakanta           | Adbhar                  |

| Schumach.                                       |                |                            |                         |
|-------------------------------------------------|----------------|----------------------------|-------------------------|
| <i>Indigofera linnaei</i> Metcalf.              | Fabaceae       | Birdsvilli indigo          | Gobra                   |
| <i>Ipomoea triloba</i> L.                       | Convolvulaceae | Littlebell                 | Gopalpur,<br>chandrapur |
| <i>Lantana camara</i> L.                        | Verbenaceae    | Red or yellow sage         | Bhudanpur               |
| <i>Leonotis nepetifolia</i> (L.) R.Br.          | Lamiaceae      | Christmas<br>Candlestick   | Ghughri chhapora        |
| <i>Lindernia crustacea</i> (L.)<br>F.Muell.     | Linderniaceae  | Brittle false<br>Pimpernel | Jetha                   |
| <i>Ludwigia perennis</i> L.                     | Onagraceae     | Evening Primrose           | Malkharauda             |
| <i>Mesosphaerum suaveolens</i> (L.)<br>Kuntze   | Lamiaceae      | Pignut                     | Adbhar                  |
| <i>Mimosa pudica</i> L.                         | Fabaceae       | Touch me not               | Hasaud                  |
| <i>Oldenlandia herbacea</i> (L.)<br>Roxb.       | Rubiaceae      | Wild coriander             | Kharsia                 |
| <i>Oxalis corniculata</i> L.                    | Oxalidaceae    | Creeping wood sorrel       | Gopalpur,<br>chandrapur |
| <i>Parthenium hysterophorus</i> L.              | Asteraceae     | Congress grass             | Saposh                  |
| <i>Persicaria lapathifolia</i> (L.)<br>Delarbre | Polygonaceae   | Pale persicaria            | Kirari                  |
| <i>Phyla nodiflora</i> (L.) Greene              | Verbenaceae    | Frogfruit                  | Chandrapur              |
| <i>Phyllanthus niruri</i> L.                    | Phyllanthaceae | Shatter stone              | Sukhada                 |
| <i>Portulaca quadrifida</i> L.                  | Portulacaceae  | Chicken weed               | Gobra                   |
| <i>Rungia muralis</i> Royle ex Nees             | Acanthaceae    | Wall rungia                | Ghughri chhapora        |
| <i>Scorzonera humilis</i> L.                    | Asteraceae     | Viper's grass              | Adbhar                  |
| <i>Senna tora</i> (L.) Roxb.                    | Fabaceae       | Sicklepod                  | Malkharauda             |
| <i>Setaria verticillata</i> (L.) P.Beauv        | Poaceae        | Bristly foxtail            | Gopalpur,<br>chandrapur |
| <i>Sida acuta</i> Burm.f.                       | Malvaceae      | Wireweed                   | Gobra                   |
| <i>Sida cordifolia</i> L.                       | Malvaceae      | Flannel weed               | Jetha                   |

|                                                           |               |                         |                      |
|-----------------------------------------------------------|---------------|-------------------------|----------------------|
| <b><i>solanum virginianum</i> L.</b>                      | Solanaceae    | Yellow nightshade fruit | Kirari               |
| <b><i>Spermacoce ocymoides</i> Burm. f.</b>               | Rubiaceae     | Basil like button weed  | Sukhada              |
| <b><i>Torenia crustacea</i> (L.) Cham. &amp; Schltdl.</b> | Linderniaceae | Malaysian false pimperl | Saposh               |
| <b><i>Trianthema portulacastrum</i> L.</b>                | Aizoaceae     | Black pigweed           | Kharsia              |
| <b><i>Tridax procumbens</i> L.</b>                        | Asteraceae    | Coat button daisy       | Chandrapur           |
| <b><i>Urochloa ramosa</i> (L.) T.Q. Nguyen</b>            | Poaceae       | Dixie signalgrass       | Gopalpur, chandrapur |
| <b><i>Xanthium strumarium</i> L.</b>                      | Asteraceae    | Rough cocklebur         | Kirari               |

## Discussion and Conclusion

The present investigation revealed a rich diversity of weed flora in the study area, with 72 species distributed across multiple taxonomic families. The dominance of families such as Asteraceae, Poaceae, Fabaceae, Amaranthaceae, and Cyperaceae is consistent with earlier floristic studies, as these families are well known for their ecological adaptability, high reproductive capacity, and efficient dispersal mechanisms. The predominance of Asteraceae and Poaceae indicates their ability to thrive under varied environmental conditions, particularly in disturbed and agricultural habitats. Similar findings have been reported in other agroecosystems, where these families contribute significantly to weed diversity and dominance.

The present study provides a comprehensive account of the weed flora occurring across different localities of the study area, highlighting considerable diversity in terms of species composition, family representation, and spatial distribution. The documentation of a wide range of weed species belonging to numerous genera

and families reflects the heterogeneous nature of habitats, including agricultural fields, fallow lands, roadsides, wastelands, and semi-natural ecosystems. Such diversity is indicative of favorable climatic conditions, variable soil types, and continuous anthropogenic disturbances that facilitate the establishment and persistence of weeds. Among the recorded taxa, families such as Asteraceae, Poaceae, Amaranthaceae, Fabaceae, and Cyperaceae were found to be dominant. The dominance of these families can be attributed to their efficient reproductive strategies, rapid growth, high seed production, and strong adaptive potential under diverse environmental conditions. Grasses and sedges, in particular, showed wide distribution across collection sites, emphasizing their ecological importance and competitive advantage in disturbed and cultivated habitats. The frequent occurrence of annual and perennial weeds further suggests their role in altering crop growth dynamics and nutrient availability. Several invasive and noxious weed species, including *Parthenium hysterophorus*, *Chromolaena odorata*, *Lantana camara*, *Cyperus rotundus*, and *Cynodon dactylon*, were commonly observed throughout the study area. The widespread presence of these species underscores their aggressive colonizing ability

and highlights the ecological and economic threats they pose to agricultural productivity and native plant diversity. Their dominance may lead to reduced crop yields, displacement of indigenous flora, and long-term degradation of soil quality if left unmanaged.

## References

1. Bell, R. S., W. H. Lachman, E. M. Rahn, and R. D. Sweet. 1962. Life History Studies as Related to Weed Control in the Northeast. I. nutgrass. Kingston, RI: Rhode Island Agricultural Experiment Station Bull. 364. 33 p.
2. Bhowmik, P. C. (1997). Weed biology: importance to weed management. *Weed science*, 45(3), 349-356.
3. Chaudhary, P., & Khalkho, A. D. (2025). A review on some traditional medicinal plants of Chhattisgarh, India. *World Wide J Multidiscip Res Dev*, 11(2), 33-36.
4. Chaudhary, P., Khalkho, A. D., & Soni, D. K. (2025). Diversity of Weed Species from Sakti District Chhattisgarh.
5. Chouhan, S., Soni, D. K., & Khalkho, A. D. (2025). Ethnobotanical documentation of medicinal plants among the Oraon tribe of Chhattisgarh.
6. Cousens, R., & Mortimer, M. (1995). *Dynamics of weed populations*. Cambridge University Press.
7. Khalkho, A. D., & Rai, A. N. (2020). Follicolous Fungi on Medicinal Plants from Tribal Area of North Chhattisgarh.
8. Naidu, V.S.G.R. (2012), Hand Book on Weed Identification Directorate of Weed Science Research, Jabalpur, India.
9. Pavlychenko, T. K. (1937). Quantitative study of the entire root system of weed and crop plants under field conditions. *Ecology* 18:62-79.
10. Pl@ntnet (2009) retrived from <https://plantnet.org/en/>.
11. Prakash, C., & Khalkho, A. D. (2025). A survey and study on ethno medicinal plant's used by tribal people in Bodla Block, Bharamdev century of kabirdham district, cg, in India.